



Distr.: General  
26 July 2006

English only



**United Nations  
Environment  
Programme**

**Stockholm Convention on Persistent Organic Pollutants  
Persistent Organic Pollutants Review Committee**

**Second meeting**

Geneva, 6–10 November 2006

Item 6 (c) of the provisional agenda\*

**Consideration of chemicals newly proposed for inclusion in  
Annexes A, B or C of the Convention: short-chained chlorinated paraffins**

**Short-chained chlorinated paraffins proposal\*\***

**Note by the Secretariat**

The annex to the present note contains a proposal by the European Union and its member States that are Parties to the Stockholm Convention on Persistent Organic Pollutants for listing short-chained chlorinated paraffins in Annexes A, B or C of the Stockholm Convention pursuant to paragraph 1 of Article 8 of the Convention. The annex is being circulated as submitted and has not been formally edited by the Secretariat.

\* UNEP/POPS/POPRC.2/1.

\*\* Stockholm Convention, Article 8, paragraph 1.

**Proposal for listing**

**Short-chained Chlorinated Paraffins**  
**(SCCPs)**

**in the Stockholm Convention**

**on Persistent Organic Pollutants**

## **Introduction**

Short-chain chlorinated paraffins (SCCPs) are a group of synthetic compounds are mainly used in metal working fluids, sealants, as flame retardants in rubbers and textiles, in leather processing and in paints and coatings.

The available data from remote areas show clearly contamination of biota and air by SCCPs. SCCPs are highly toxic to aquatic organisms. They do not break down naturally and tend to accumulate in biota. Their persistence, bioaccumulation, potential for long-range environmental transport and toxicity mean that they may have damaging environmental effects at a global level.

This dossier focuses solely on the information required under paragraphs 1 and 2 of Annex D of the Stockholm Convention and it is mainly based on the extensive EU Risk Assessment Report on SCCPs (European Commission 2000), which is publicly available at: <http://ecb.jrc.it/existing-chemicals/>. That report and its draft update (European Commission 2005) serve as important sources of detailed technical information on SCCPs.

A large amount of other relevant summary reports and documents has been published on SCCPs in recent years and can be of value in the POP review. First and foremost, the final draft dossier prepared by Canada (Filyk et al. 2003) is a comprehensive review of highly relevant information and should thus be used in parallel to this dossier. There are also other important summary reports such as the Environmental Health Criteria report on chlorinated paraffins (WHO 1996).

## **1 Chemical identity**

SCCPs are n-paraffins that have a carbon chain length of between 10 and 13 carbon atoms and a degree of chlorination of more than 48% by weight. There is a range of commercially available C10-13 chlorinated paraffins and they are usually mixtures of different carbon chain lengths and different degrees of chlorination although all have a common structure in that no secondary carbon atom carries more than one chlorine.

Two other groups of chlorinated paraffins are made commercially. These are known as “mid, medium or intermediate chain length” (typically C14-17) and “long chain length” (typically C20-30). This dossier, however, concerns only the short chain length (C10-13) chlorinated paraffins.

### **1.1 Names and registry numbers**

IUPAC Name: Alkanes, C<sub>10-13</sub>, chloro

CAS No: 85535-84-8


EINECS No: 287-476-5

Synonyms: alkanes, chlorinated; alkanes (C<sub>10-13</sub>), chloro-(50-70%); alkanes (C<sub>10-13</sub>), chloro-(60%); chlorinated alkanes, chlorinated paraffins; chloroalkanes; chlorocarbons; polychlorinated alkanes; paraffins chlorinated.

## 1.2 Structure

Molecular formula:  $C_xH_{(2x-y+2)}Cl_y$ , where  $x=10-13$  and  $y=1-13$

The structure of two example SCCP compounds ( $C_{10}H_{17}Cl_5$  and  $C_{13}H_{22}Cl_6$ ) is shown below.



Molecular weight: 320-500

## 2 Persistence

Atmospheric half-lives of 1.9-7.2 days have been estimated for SCCPs (European Commission, 2000). Nevertheless, in the aqueous phase, rates of hydrolysis, photolysis with visible or near UV radiation, oxidation and volatilization are insignificant under ambient temperatures (Government of Canada 1993). Photolytical degradation in aqueous media may take place, but at a very slow rate (Koh and Thiemann, 2001, El-Morsi et al. 2002)

SCCPs are not readily or inherently biodegradable in standard tests. It can be concluded from the simulation tests that SCCPs with low chlorine content (e.g. <50% wt Cl) may biodegrade slowly in the environment in the presence of adapted microorganisms. An additional simulation study on degradation of SCCPs in marine environment has been required recently within the EU and the results of the study should be available by the end of 2006.

In its opinion given in 2003 (CSTEE 2003), the EU Scientific Committee on Toxicity, Ecotoxicity and the Environment came to the conclusion that SCCPs are potentially persistent (P) and possibly very persistent (vP). The Committee emphasised the evidence that SCCPs are occurring in remote areas and were of the view that this is particularly important evidence that gives further support to the P/vP classification. Weight of evidence indicates that the half-life of SCCPs in sediment is greater than 1 year

As the available information seems to indicate that SCCPs have long half-lives in the environment, it can be concluded that SCCPs meet the screening criteria for persistency laid down in Annex D of the Stockholm Convention.

## 3 Bioaccumulation

Reported log Kow of different SCCPs range from 4.39 to 8.69, indicating a high potential for bioaccumulation (European Commission, 2000).

High bioconcentration factors in fish have been reported in the scientific literature (European Commission, 2000). In one of the key studies, whole body bioconcentration factors (BCFs) of 1

173-7 816 were determined based on radioactivity measurements in the fish and BCFs of 574-7 273 were determined based on the parent compound analysis (Madeley and Maddock 1983).

In addition to these experimental values from laboratory studies, BCF values have been estimated *in situ* for lake trout (*Salvelinus namaycush*) in western Lake Ontario. The overall BCF for SCCPs (C<sub>10-13</sub>) in lake trout from western Lake Ontario was 36 500 (Filyk et al. 2003). Bioconcentration in mussels has also been assessed with reported whole body BCFs ranging from 5 785 to 40 900 (European Commission 2000).

Therefore, in view of the high reported values of both log K<sub>ow</sub> and BCF for different aquatic species, it can be concluded that SCCPs meet the screening criteria for bioaccumulation laid down in Annex D of the Stockholm Convention.

#### 4 Potential for long-range environmental transport

Drouillard et al. (1998a) have determined vapour pressures for a range of SCCPs. In the EU Risk Assessment Report (European Commission, 2000), an assumed vapour pressure of a SCCP with chlorine content of approximately 50% of 0.0213 Pa at 40°C is used. Henry's law constants range from 0.7 to 18 Pa m<sup>3</sup>/mol, near the values of some acknowledged POPs (Drouillard et al 1998a). This constant and vapour pressure are the most important chemical characteristic to determine whether a substance may undergo long-range environmental transport in the atmosphere. As shown in table 1, Henry's law constants are in the range of those for currently listed POPs. Therefore, attending to their chemical properties, SCCP are likely to undergo long range environmental transport in the atmosphere. Moreover, atmospheric half-lives exceeding the screening criteria of 2 days (1.9-7.2 days) have been estimated for SCCPs (European Commission, 2000).

**Table 1: Water solubility (WS), vapour pressure (VP) and Henry's Law Constant (HLC) (at 25 °C) for SCCPs and currently listed POPs.**

Substance	WS mg/L	VP Pa	HLC Pa m <sup>3</sup> /mol
SCCP-min	0.0224**	2.8 x 10 <sup>-7</sup> *	0.7 *
SCCP-max	0.994**	2.5 *	18 *
POP-min	0.0012 (DDT)	2.5 x 10 <sup>-5</sup> (DDT)	0.04 (endrin)
POP-max	3.0 (toxaphene)	27 (toxaphene)	3726 (toxaphene)
POP-2 <sup>nd</sup> max	0.5 (dieldrin)	0.04 (heptachlor)	267 (heptachlor)

\* Drouillard et al. 1998a

\*\* Drouillard et al. 1998b

The levels of SCCPs in air have been found in samples from a remote area in the Canadian Arctic (Peters et al., 1998). The mean total (vapour + particulate phase) levels found were 20±32 pg/m<sup>3</sup> at the remote site. Tomy (1997a; as reported in Tomy, 1998) found that SCCPs were present in air from Egbert, Canada at a concentration of 65-924 pg/m<sup>3</sup>. Muir et al. (2001) reported short chain chlorinated paraffins to be present at a concentration of 249 pg/m<sup>3</sup> in air overlying the west basin of Lake Ontario. The levels of SCCPs in air from the Arctic have also been reported by Bidleman et al. (2001). The levels found ranged from 1.07 to 7.25 pg/m<sup>3</sup> and were dominated by the contributions from chlorodecanes (C<sub>10</sub> fractions).

The levels of SCCPs (vapour + particulate phase) reported in the Arctic environment ranged from 1.07 to 7.25 pg/m<sup>3</sup> (Borgen et al., 2000) and from 1.8 to 10.6 ng/m<sup>3</sup> (Bidleman et al., 2001). Also Tomy et al. (1997 and 1999) reported occurrence of SCCPs in surface sediments from the Canadian mid-latitude and Arctic regions, attributed to long-range transport. Stern (2003, as reported in Filyk, 2003) found levels of SCCPs in a lake sediment core taken from a very remote lake in the Arctic (75°34'N; 89°19'W) which provide evidence for transport to and deposition in the Arctic (Filyk et al. 2003).

Levels of SCCPs in marine mammals from various regions of the Arctic have been reported (Stern et al., 1997), as well as from Canada and Greenland (Tomy et al., 1998). There is also evidence of SCCPs accumulation in fish species from Lake Ontario (Muir et al., 2001).

The study of Stern et al. (1997) also detected SCCPs in three samples of breast milk taken from Inuit women living in settlements along the Hudson Strait. A study performed by Thomas et al. (2006) found similar SCCP concentrations in breast milk of women from United Kingdom.

The ubiquity of SCCP and the vapour pressure values that have been reported, together with the Henry's constant values (similar to those of acknowledged POPs), give enough evidence to conclude that SCCPs meet the screening criteria for long range environmental transport laid down in Annex D of the Stockholm Convention..

## **5 Adverse effects**

According to EU Risk Assessment Report (European Commission, 2000) SCCPs are of low acute toxicity to fish with 48 - and 96-hour LC<sub>50</sub>s in excess of the water solubility of the substance. Chronic toxicity values include a 60-day LC<sub>50</sub> of 0.34 mg/l for rainbow trout and no observed effect concentrations of <0.040 and 0.28 mg/l for rainbow trout and sheepshead minnow respectively.

For aquatic invertebrates, SCCPs are of high toxicity with 24-hour EC<sub>50</sub>s with daphnids ranging from 0.3 to 11.1 mg/l and with acute NOECs ranging from 0.06 to 2 mg/l. In 21-day tests with daphnids, EC<sub>50</sub>s ranged from 0.101 to 0.228 mg/l and NOECs ranged from 0.005-0.05 mg/l. For algae, 96-hour EC<sub>50</sub>s ranged from 0.012 to 3.7 mg/l, depending on the species.

Information available from acute studies and skin irritation studies in animals indicates that the intensity and nature of effects for these endpoints are independent of chain length and degree of chlorination. Assessment of the available data clearly indicates that SCCPs are of low acute toxicity in animals. In rodent carcinogenicity studies, dose-related increases in the incidence of adenomas and carcinomas were observed in the liver, thyroid and kidney. Other cancers seen were dismissed as not significant. Consideration of the likely underlying mechanisms for these tumours suggests that they are not relevant to human health.

There are no data available in humans or animals on fertility. A SCCP produced developmental effects in rats at a dose which also caused maternal toxicity (2 000 mg/kg). No developmental effects were observed in a study in rabbits, although maternally toxic doses were not tested. NOAELs for general toxicity of 100 and 1 000 mg/kg/day were identified in rats and mice respectively.

The relevance of the finding that medium-chain length chlorinated paraffins can cause a severe effect (internal haemorrhaging leading to deaths) in suckling rat pups has been recently discussed (European Commission, 2005).

The current EU hazard classification for SCCPs is: Carc. Cat. 3; R40 - N; R50-53 (Risk Phrases: R40: Limited evidence of a carcinogenic effect; R50/53: Very toxic to aquatic organisms; may cause long-term adverse effects in the aquatic environment.). Also the International Agency for Research on Cancer (IARC) has designated SCCPs (as a group) as possible carcinogens.

To sum up, SCCPs are of high aquatic toxicity to a variety of animals, and its terrestrial toxicity may be an additional cause of concern. As it's also a potential carcinogen, it fulfils the screening criteria for adverse effects to human health and the environment.

## **6 Statement of the reasons for concern**

SCCPs are highly toxic to aquatic organisms and it is considered as a possible carcinogen. SCCPs do not break down naturally and tend to accumulate to biota. The available data from remote areas show clearly contamination of environment and biota by SCCPs. Their persistence, bioaccumulation and toxicity mean that they may have damaging environmental effects at a global level. Overall, it can be considered that SCCPs meet the screening criteria for persistence, potential to cause adverse effects, bioaccumulation and potential for long range environmental transport.

Placing on the market and use of SCCPs has been restricted over the last years in the European Union but no total prohibition has yet been foreseen. On the other hand, production and use of SCCPs continues unrestricted in many other countries. As SCCPs can move in the atmosphere far from its sources, single countries or groups of countries alone cannot abate the pollution caused by it. Due to the harmful POP properties and risks related to its widespread production and use, international action is warranted to control this pollution.

## **7 References**

Bidleman T. F., Alaee M. and Stern G. A. (2001). New persistent chemicals in the Arctic environment. In Synopsis of research conducted under the 2000-2001 Northern Contaminants Program, Edited by S. Kalhok. - Ottawa : Dept. of Indian Affairs and Northern Development, 2001, p. 93-103, ill. Abstract available from [www.inac.gc.ca/ncp](http://www.inac.gc.ca/ncp).

Borgen A. R., Schlabach M. and Gundersen H. (2000). Polychlorinated alkanes in arctic air. *Organohalogen Compounds*, 47, 272-275.

CSTEE (2003). Opinion of the Scientific Committee on Toxicity, Ecotoxicity and the Environment on the scientific basis of the national provisions on short chain chlorinated paraffins (SCCPs) being more restrictive than those laid down in directive 2002/45/EC that the Netherlands intends to maintain in accordance with Article 95(4) of the EC Treaty. Adopted by the CSTEE by written procedure on 3 October 2003, [http://europa.eu.int/comm/health/ph\\_risk/committees/sct/documents/out200\\_en.pdf](http://europa.eu.int/comm/health/ph_risk/committees/sct/documents/out200_en.pdf).

Drouillard K. G., Tomy G. T., Muir D. C. G. and Friesen K. J. (1998a). Volatility of chlorinated n-alkanes (C10-C12): Vapour pressures and Henry's Law constants. *Environmental Toxicol. Chem.*, 17, 1252-1260.

Drouillard, K.G.; Heibert, T.; Tran, P.; Tomy, G.T.; Muir, D.C.G.; Friesen, K.J. (1998b) Estimating the aqueous solubilities of individual chlorinated n-alkanes (C10-C12) from measurements of chlorinated alkane mixtures. *Environ. Toxicol. Chem.* 17, 1261-1267.

El-Morsi, T.M., Emar, M.M.; El Bary, H.M.H.A., Abd-El-Aziz, A.S., and Friesen, K.J. (2002). Homogeneous degradation of 1,2,9,10-tetrachlorodecane in aqueous solutions using hydrogen peroxide, iron and UV light. *Chemosphere* 47:343-348

European Commission (2000). European Union Risk Assessment Report. Vol. 4: alkanes, C10–13, chloro-. European Chemicals Bureau, Brussels, Belgium. 166 pp. (EUR 19010; ISBN 92-828-8451-1).

European Commission (2005). Final Draft. Updated Risk Assessment of alkanes, C10–13, chloro-. Combined draft of August 2005.

Filyk, G, Lander, L, Eggleton, M, Muir, D, Puckett, K. (2003) Short Chain Chlorinated. Paraffins (SCCP) Substance-Final Draft II. Environment Canada. Dossier prepared for UNECE ad hoc Expert Group on POPs. Available at: <http://www.unece.org/env/popsxg>

Government of Canada (1993). Priority Substances List Assessment Report. Chlorinated paraffins. Minister of Supply and Services, Ottawa, Ontario (ISBN 0-662-20515-4; Catalogue No. En40-215/17E). 32 pp.

Koh, I.-O., and Thiemann, W.H.-P. (2001). Study of photochemical oxidation of standard chlorinated paraffins and identification of degradation products. *J Photochem. Photobiol. A* 139: 205-215.

Madeley J. R. and Maddock B. G. (1983). The bioconcentration of a chlorinated paraffin in the tissues and organs of rainbow trout (*Salmo gairdneri*). ICI Confidential Report BL/B/2310.

Muir D., Bennie D., Teixeira C., Fisk A., Tomy G., Stern G. and Whittle M. (2001). Short chain chlorinated paraffins: Are they persistent and bioaccumulative? ACS Symposium Series, 773, 184-202.

Peters A. J., Tomy G. T., Stern A. S. and Jones C. J. (1998). Polychlorinated alkanes in the atmosphere of the United Kingdom and Canada – Analytical methodology and evidence of the potential for long-range transport. *Organohalogen Compounds*, 35, 439-442.

Stern G. A., Tomy G. T., Muir D. C. G., Westmore J. B., Dewailly E., and Rosenberg B. (1997). Polychlorinated n-alkanes in aquatic biota and human milk. Convention on Long-range Transboundary Air Pollution. Working Group on Strategies (Twenty-first session, 16-20 June 1997). Informed in-session document No.2.

Stern G. A. (2003) Sediments. In Canadian Arctic Contaminants Assessment Report, Phase 2. Bidleman, T.F., Macdonald, R.W., Stow, J. (eds.) Department of Indian Affairs and Northern Development, Ministry of Public Works and Government Services Canada, QS-8525-002-EE-A1, R72-260/2003-2E, 0-662-33468X, 332 pages.

Thomas, G.O., Farrar, D., Braekevelt, E., Stern, G., Kalantzi, O.I., Martin, F.L.; and Jones, K.C. (2006). Short and medium chain length chlorinated paraffins in UK human milk fat *Environment International* 32:34-40



Tomy G. T. (1997a). The mass spectrometric characterization of polychlorinated n-alkanes and the methodology for their analysis in the environment. PhD Thesis, University of Manitoba, Winnipeg (as quoted in Tomy, 1998).

Tomy G. T., Stern G. A., Muir D. C. G., Lockhart L. and Westmore J. B. (1997b). Occurrence of polychloro-n-alkanes in Canadian mid-latitude and Arctic lake sediments. *Organohalogen Compounds*, 33, 220-226.

Tomy G. T. (1998). Environmental chemistry and toxicology of polychlorinated n-alkanes. *Rev. Environ. Contam. Toxicol.*, **158**, 53-128.

Tomy G. T., Stern G. A., Lockhart W. L. and Muir D. C. G. (1999). Occurrence of C10-C13 polychlorinated n-alkanes in Canadian midlatitude and Arctic lake sediments. *Environ. Sci. Technol.*, 33, 2858-2863.

---